

One-Dimensional Kondo Lattice at Half Filling: Charge Properties.

A. M. Tsvelik

February 1, 2008

*Department of Physics, University of Oxford, 1 Keble Road,
Oxford, OX1 3NP, UK*

Abstract

The model of Kondo chain with M -fold degenerate band of conduction electrons of spin $1/2$ at half filling interacting with localized spins S is studied. It is shown that due to the instanton effects the spin sector reveals itself in optical conductivity. Namely, the density of the charge current at large wave vector $k \approx \pi$ is proportional to the density of topological charge of the staggered magnetization. As a consequence, the optical conductivity at $k = \pi$ would either have a gap equal to three spin gaps (for $|M - 2S| = (\text{even})$ when the spin spectrum is gapful), or show a pseudogap $\sigma(\omega, k = \pi) \sim \omega^2$ for $|M - 2S| = (\text{odd})$.

PACS numbers: 74.65.+n, 75.10. Jm, 75.25.+z

In this letter I continue to study the model of one dimensional of Kondo chain at half filling introduced in the previous publication ¹. This model has the following Hamiltonian:

$$H = \sum_r \sum_{\alpha=1}^M \left[-\frac{1}{2} (c_{r+1,\alpha,a}^+ c_{r,\alpha,a} + c_{r,\alpha,a}^+ c_{r+1,\alpha,a}) + J (c_{r,\alpha,a}^+ \hat{\sigma}_{ab} c_{r,\alpha,b}) \vec{S}_r \right] \quad (1)$$

The conduction electrons have spin $1/2$ and their band has an additional M -fold degeneracy. The value of local spins is S .

It was shown in Ref. 1 that the low energy behavior of the Kondo chain with a half filled electron band is determined by antiferromagnetic correlations (the

RKKY coupling) and the Kondo effect plays no role. Depending on the value of $2S - M$ the spin excitations are either gapless (if $2S - M$ is odd) or have a spectral gap exponentially small in t/J ($2S - M$ is even). Thus the spin correlation length is always large and the low energy sector can be described by a continuous field theory. It was shown that this theory is the $O(3)$ nonlinear sigma model with the topological term. The coefficient of the topological term is $\pi(\text{mod } 2\pi)$ for odd $2S - M$ and $0 \pmod{2\pi}$ for even $2S - M$. These results are in qualitative agreement with the strong coupling picture and with the numerical results of other authors^{2,3}.

In the previous publication I made some remarks about the charge sector of the model. In particular I argued that the charge gap was equal to the spin gap and provided the explicit expression for the wave function of an electron-kink bound state. These arguments were justly criticized by X.-G. Wen who pointed out that the suggested wave function is not normalizable⁴.

In this letter I present rigorous results for the charge sector of the model (1). It turns out that despite of the fact that the charge excitations do have a large gap $\Delta_{ch} \approx JS$, the spin sector contributes into the optical conductivity at $k = \pi$ through instantons. This contribution is purely dynamical. For even $2S - M$ the indirect gap measured by optical conductivity is equal to $3\Delta_{sp}$ (Δ_{sp} is the spin gap) and for odd $2S - M$ there is only a pseudogap.

In order to analyse charge properties of the model (1), we have to recall some arguments of Ref. 1. The suggested approach is based on the fact that at $J/t \ll 1$ the spin subsystem is very slow. This allows to treat the staggered magnetization $\vec{n}$ as a slow variable and linearize the fermionic spectrum around the Fermi points

$$\begin{aligned}\vec{m}_r &= a\vec{k}(x) + (-1)^r \vec{n}(x) \sqrt{1 - a^2 \vec{k}(x)^2}, (\vec{k}\vec{n}) = 0, \\ c_r &= i^r \psi_R(x) + (-i)^r \psi_L(x)\end{aligned}\tag{2}$$

Substituting this expression into the initial model (1) we get the following Lagrangian density for fermions:

$$L = \bar{\psi}_j [i\gamma_\mu \partial_\mu \hat{I} + JS(\hat{\sigma}\vec{n}(x))\sqrt{1 - a^2 \vec{k}(x)^2}] \psi_j\tag{3}$$

where the term coupled to $\vec{k}$ is omitted as irrelevant. The corresponding fermionic determinant was calculated in Ref. 1. In what follows we shall be interested in the expansion of this determinant in gradients of $\vec{n}$. The coordinate dependence of the amplitude of the mass term coming from the $\sqrt{1 - a^2 \vec{k}(x)^2}$ -term is not

important in this respect. Therefore we shall treat this term as constant: $JS < \sqrt{1 - a^2 \vec{k}(x)^2} = m$. It was shown that this determinant was a particular case of the more general one:

$$D[g] = MTr \ln[i\gamma_\mu \partial_\mu + (1 + i\gamma_S)mg/2 + (1 - i\gamma_S)mg^+/2] = MW(g) \quad (4)$$

$$W(g) = \int d^2x \left\{ \frac{1}{16\pi} Tr(\partial_\mu g^+ \partial_\mu g) + \frac{1}{24\pi} \int_0^1 d\xi \epsilon_{abc} Tr(g^+ \partial_a g g^+ \partial_b g g^+ \partial_c g) \right\} \quad (5)$$

where g belongs to the SU(2) group, $\gamma_S = i\sigma_3$ and $W(g)$ is the famous Wess-Zumino-Witten action. The equivalence becomes obvious when one makes the transformation

$$\psi_R \rightarrow i\psi_R, \psi_L \rightarrow \psi_L \quad (6)$$

and put $n_0 = 0$ in the general expression for the matrix g :

$$g = n_0 + i(\vec{\sigma}\vec{n})\sqrt{1 - n_0^2} \quad (7)$$

The derivation of Eq.(4) given in Ref. 1. can be simplified. Namely, one can use the procedure of non-Abelian bosonization (see Refs. 5 - 7, I also recommend the book by Itzykson and Drouffe⁸). In this procedure the action of free fermions on a group U(2) (this can be done for any group, but I do not want to complicate the arguments) can be rewritten as follows:

$$\int d^2x i\bar{\psi}_a \gamma_\mu \partial_\mu \psi_a = W(h) + \frac{1}{4\pi} \int d^2x (\partial_\mu \phi)^2, \quad (8)$$

$$\psi_{R,a}^+ \psi_{L,b} \sim e^{i\phi} h_{ab} \quad (9)$$

where h belongs to the SU(2) group and W is given by Eq.(refeq:Det). Using this equivalence we get the following:

$$\begin{aligned} S &= \int d^2x [i\bar{\psi}_a \gamma_\mu \partial_\mu \psi_a + \frac{m}{2} \bar{\psi}_a (1 + i\gamma_S) g_{ab} \psi_a + \\ &\quad \frac{m}{2} \bar{\psi}_a (1 - i\gamma_S) g_{ab}^+ \psi_b] = \\ &\frac{1}{4\pi} \int d^2x (\partial_\mu \phi)^2 + W(h) + \tilde{m} \int d^2x Tr(e^{i\phi} g^+ h + H.c) \end{aligned} \quad (10)$$

Now we do the transformation $g^+ h = G$ and use the well-known identity

$$W(gG) = W(G^+) + W(g) + \frac{1}{16\pi} \int d^2x (g^+ \partial_{\bar{z}} g G \partial_z G^+) \quad (11)$$

As a result we get

$$S = S_1 + W(g) \quad (12)$$

$$S_1 = \frac{1}{4\pi} \int d^2x (\partial_\mu \phi)^2 + W(G^+) + \tilde{m} \int d^2x Tr(e^{i\phi} G + H.c) \quad (13)$$

Since the operator $e^{i\phi}G$ has the right and left conformal dimensions $(1/2, 1/2)$, it is relevant and the theory described by the action S_1 is massive with the mass gap $\tilde{m} \approx JS$. The remaining part of the action is equal to $W(g)$ which reproduces the result Eq.(4).

Now we shall use the identity (4) to calculate the correlation function of electronic currents. For this purpose we calculate the variational derivative of Eq.(4) with respect to n_0 at $n_0 = 0$ to establish the following Ward identity:

$$J_S = m\bar{\psi}\gamma_S\psi = \frac{\delta D(g)}{\delta n_0}|_{n_0=0} = -\frac{M}{4\pi}\frac{\delta}{\delta n_0} \int d^2x Tr[\delta g g^+ \partial_\mu(g^+ \partial_\mu g)] = \frac{iM}{\pi}\epsilon_{\mu\nu}(\vec{n}[\partial_\mu \vec{n} \times \partial_\nu \vec{n}]) \quad (14)$$

Thus the anomalous fermionic current is proportional to the density of topological charge of the staggered magnetization. In this way the charge and the spin sector are related to each other.

In terms of the original fermions the anomalous current is nothing but the ordinary current at $k = \pi$. Therefore the pair correlation function of J_S is proportional to the optical conductivity at $k = \pi$:

$$\sigma(\omega, k = \pi - q) = \frac{e^2}{\omega} \Im m << J_S(-\omega, -q) J_S(\omega, q) >> \quad (15)$$

In the case $2S - M = (\text{even})$ the spin sector is described by the ordinary O(3) nonlinear sigma model. The spin sector has a gap and the correlation function of topological densities decay exponentially in Euclidean space-time. It was shown by Kirillov and Smirnov⁹ that the first non-vanishing formfactor in the Lehmann expansion of this correlation function contains three elementary excitations, that is the gap measured by the optical conductivity is equal to $3\Delta_{sp}$. The case $2S - M = (\text{odd})$ is more interesting. In this case the spin sector is critical and is described by the O(3) nonlinear sigma model with $\theta = \pi$. The latter is equivalent to the $S = 1/2$ Heisenberg chain, that is can be written as a theory of the massless scalar bosonic field Φ :

$$S_\pi = \frac{1}{2} \int d^2x (\partial_\mu \Phi)^2 \quad (16)$$

The components of the staggered magnetization are expressed as exponents of Φ and the dual field Θ defined as

$$\epsilon_{\mu\nu} \partial_\nu \Theta = \partial_\mu \Phi \quad (17)$$

namely¹⁰,

$$\vec{n} \sim (\cos(\sqrt{2\pi}\Phi), \cos(\sqrt{2\pi}\Theta), \sin(\sqrt{2\pi}\Theta)) \quad (18)$$

Substituting this expression into Eq.(14) and using the identity (17) we get

$$J_S \sim \sin(\sqrt{2\pi}\Phi)(\partial_\mu\Phi)^2 \quad (19)$$

The pair correlation function of the currents is

$$<< J_S(-\omega_n, -q) J_S(\omega_n, q) >> \sim (\omega_n^2 + v^2 q^2)^{3/2} \quad (20)$$

and the optical conductivity

$$\sigma(\omega, k = \pi - q) \sim \frac{1}{\omega} \Im m[\omega^2 - v^2 q^2]^{3/2} \quad (21)$$

where v is the velocity of spin waves. The latter means that the optical conductivity would reveal in this case only a pseudogap at $k = \pi$.

As we see, from one side it is fair to say that the spectrum of the system is separated into two sectors (we call them the charge and the spin sectors) and the charge sector always has a large gap Δ_{ch} of order of JS in agreement with the strong coupling expansion and the results of Refs. 2, 3. This conclusion follows from the discussion around Eq.(13). It does not mean however, that optical measurements measure only Δ_{ch} (they do it only at small wave vectors). Due to the fact that the physical fermions are composite fields, there is the admixture of spin sector. The very form of the identity (14) tells us that this admixture originates from instantons, which is not unusual¹⁰.

I am grateful to X.-G. Wen and M. Sigrist for the constructive criticism, for I. Lerner and to A. Ludwig for the interest to the work.

References

- [1] A. M. Tsvelik, Phys. Rev. Lett. **72**, 1048 (1994).
- [2] H. Tsunetsugu, Y. Hatsugai, K. Ueda and M. Sigrist, Phys. Rev. **B 46**, 3175 (1992)
- [3] C. C. Yu, S. R. White, Phys. Rev. Lett. **71**, 3866 (1993).
- [4] X.-G. Wen, private communication.
- [5] E. Witten, Comm. Math. Phys. **92**, 455 (1984).
- [6] A. M. Polyakov and P. B. Wiegmann, Phys. Lett. **B 131**, 121 (1983).

- [7] V. G. Knizhnik and A. B. Zamolodchikov, Nucl. Phys. **B247**, 83 (1984).
- [8] C. Itzykson and J. - M. Drouffe, in *Statistical Field Theory*, v. 2, Appendix 9.C, Cambridge University Press Publ. (1989).
- [9] A. N. Kirillov and F. A. Smirnov, Int. J. Mod. Phys. **3**, 731 (1988).
- [10] For a review see chapters about instantons in S. Coleman *Aspects of Symmetry*, Cambridge University Press, (1988).